

Vol 3 Issue 9 Oct 2013

ISSN No : 2230-7850

---

Monthly Multidisciplinary  
Research Journal

*Indian Streams  
Research Journal*

Executive Editor

Ashok Yakkaldevi

Editor-in-chief

H.N.Jagtap

---

## Welcome to ISRJ

**RNI MAHMUL/2011/38595**

**ISSN No.2230-7850**

Indian Streams Research Journal is a multidisciplinary research journal, published monthly in English, Hindi & Marathi Language. All research papers submitted to the journal will be double - blind peer reviewed referred by members of the editorial Board readers will include investigator in universities, research institutes government and industry with research interest in the general subjects.

### **International Advisory Board**


Flávio de São Pedro Filho Federal University of Rondonia, Brazil	Mohammad Hailat Dept. of Mathematical Sciences, University of South Carolina Aiken, Aiken SC 29801	Hasan Baktir English Language and Literature Department, Kayseri
Kamani Perera Regional Centre For Strategic Studies, Sri Lanka	Abdullah Sabbagh Engineering Studies, Sydney	Ghayoor Abbas Chotana Department of Chemistry, Lahore University of Management Sciences [ PK ]
Janaki Sinnasamy Librarian, University of Malaya [ Malaysia ]	Catalina Neculai University of Coventry, UK	Anna Maria Constantinovici AL. I. Cuza University, Romania
Romona Mihaila Spiru Haret University, Romania	Ecaterina Patrascu Spiru Haret University, Bucharest	Horia Patrascu Spiru Haret University, Bucharest, Romania
Delia Serbescu Spiru Haret University, Bucharest, Romania	Loredana Bosca Spiru Haret University, Romania	Ilie Pinteau, Spiru Haret University, Romania
Anurag Misra DBS College, Kanpur	Fabricio Moraes de Almeida Federal University of Rondonia, Brazil	Xiaohua Yang PhD, USA Nawab Ali Khan College of Business Administration
Titus Pop	George - Calin SERITAN Postdoctoral Researcher	

### **Editorial Board**

Pratap Vyamktrao Naikwade ASP College Devrukh,Ratnagiri,MS India	Iresh Swami Ex - VC. Solapur University, Solapur	Rajendra Shendge Director, B.C.U.D. Solapur University, Solapur
R. R. Patil Head Geology Department Solapur University, Solapur	N.S. Dhaygude Ex. Prin. Dayanand College, Solapur	R. R. Yaliker Director Managment Institute, Solapur
Rama Bhosale Prin. and Jt. Director Higher Education, Panvel	Narendra Kadu Jt. Director Higher Education, Pune	Umesh Rajderkar Head Humanities & Social Science YCMOU, Nashik
Salve R. N. Department of Sociology, Shivaji University, Kolhapur	K. M. Bhandarkar Praful Patel College of Education, Gondia	S. R. Pandya Head Education Dept. Mumbai University, Mumbai
Govind P. Shinde Bharati Vidyapeeth School of Distance Education Center, Navi Mumbai	Sonal Singh Vikram University, Ujjain	Alka Darshan Shrivastava Shaskiya Snatkottar Mahavidyalaya, Dhar
Chakane Sanjay Dnyaneshwar Arts, Science & Commerce College, Indapur, Pune	G. P. Patankar S. D. M. Degree College, Honavar, Karnataka	Rahul Shriram Sudke Devi Ahilya Vishwavidyalaya, Indore
Awadhesh Kumar Shirotriya Secretary, Play India Play (Trust),Meerut	Maj. S. Bakhtiar Choudhary Director,Hyderabad AP India.	S.KANNAN Ph.D , Annamalai University,TN
	S.Parvathi Devi Ph.D.-University of Allahabad	Satish Kumar Kalhotra
	Sonal Singh	

**Address:-Ashok Yakkaldevi 258/34, Raviwar Peth, Solapur - 413 005 Maharashtra, India  
Cell : 9595 359 435, Ph No: 02172372010 Email: ayisrj@yahoo.in Website: www.isrj.net**



 **SEASONAL VARIATION IN DIATOMS DENSITY AND THEIR CORRELATION WITH PHYSICO-CHEMICAL PARAMETERS OF YASHWANT LAKE, TORANMAL (M.S.) INDIA**



Ekhande A. P<sup>1</sup>, Patil J. V<sup>1</sup> And Padate G.<sup>2</sup> S

<sup>1</sup>SVS's Dadasaheb Rawal College, Dondaicha, Dist. Dhule (M.S.) India.

<sup>2</sup>Dept. of Zoology, Faculty of Science, The M. S. University of Baroda, Vadodara

**Abstract:** Seasonal variation of diatoms density and species richness was studied for two years from Yashwant Lake, located in the Satpura Reserve Forest at Toranmal (Maharashtra) India. The density of diatoms was minimal in post-monsoon which increased gradually and culminated in summer. Species richness of diatoms was maximal in summer and found to be minimal in winter. The Pearson correlation was calculated by keeping phytoplankton as dependent variable and other abiotic factors as independent variables. The diatoms structure depends on a variety of environmental factors that include biological parameters as well as various physico-chemical factors. The Yashwant Lake, located in subtropics, receives maximum photoperiod during summer that invigorates growth of the diatoms. In post monsoon the water cover and water level were higher which distribute the phytoplankton and leads to decrease in their density, while opposing situations occurs in summer leading to increase in density of total phytoplankton.

**Keywords:** Toranmal, Diatoms, Seasonal variation, density, species richness

#### INTRODUCTION

The Phytoplankton of the open water ponds, lakes and large streams consist of a diverse assemblage of microscopic autotrophs. Many of them have different physiological requirements, which may vary in response to physical and chemical parameters such as light, temperature and nutritional regime (Wetzel, 2001). The dominant genera in algal groups not only change spatially (vertically and horizontally within a lake) but also show seasonal variations in response to seasonal changes in physical, chemical and biological conditions of the water body. Hence a general pattern of seasonal succession of phytoplankton of many lakes has been correlated with environmental factors. The precise reasons for many of these changes are not well known (Wetzel, 2001).

The studies on phytoplankton distribution are pivotal in understanding the health of particular water body. They are important source of fish diet, as pollution indicator and they project the trophic status of the water body (Naik and Neelkanthan, 1990). This heterogeneous microscopic group include members of different families such as Cyanophyceae, Chlorophyceae, Bacillariophyceae (diatom), Dinophyceae and Euglenophyta.

At present, altogether fortyone functional freshwater phytoplankton groups are described worldwide with more or less precisely defined ecological demands (Padisak et al, 2009). Further, field and laboratory evidences indicate that phytoplankton react rapidly not only to the climate but also to the changes in nutrient loading.

Therefore, they either mimic eutrophication by increasing phytoplankton production carrying capacity of ecosystem (Mooij et al, 2005) or contribute to it by inducing phosphorus release from sediments and accelerating the nutrient cycling (Pettersson et al, 2003).

The diatoms in Littoral zone are important contributors of the primary productivity in shallow aquatic ecosystems (Wetzel, 1990). According to Palmer (Palmer, 1980) *Ulnaria acus*, *Gomphonema sp.*, *Cyclotella sp.* and *Melosira sp.* are found in organically rich water and play an important role in water quality assessment and trophic structure. Diatoms are important in Paleolimnological studies to reconstruct the past eutrophication of lakes on basis of paleolimnological evidences (Taylor et al, 2006).

In the present study, an attempt is made to find out the effect of season on the distribution of phytoplankton in Yashwant Lake that receives the southwest monsoon and has altitudinal effect.

#### MATERIAL AND METHODS

Yashwant Lake is a perennial water body, located on Toranmal Plateau, one of the important plateaus in mid Satpura. This plateau forms a table land on the summit, covering about 41 Sq. Km. area at 1155 meter altitude (AMSL) extending between 21° 54' North to 21° 61' latitude and 74° 26' to 74° 34' East longitude. Yashwant Lake has a perimeter of 2.75 Km. and spreads in 39 hectares. It was constructed during British period by damming the dip

gorge. The littoral zone of Yashwant Lake is covered with various macrophytes. The Lake on the west and northwest is surrounded by forested land.



Google Sat. Image of Yashwant Lake

The diatoms along the periphery of Yashwant Lake were collected after the interval of every fifteen days from three stations namely Yashwant Lake –A-site (YLA), Yashwant Lake –B-site (YLB) and Yashwant Lake –C-site (YLC). Ten liters of water was filtered through the plankton net No. 25 of bolting silk with mesh size 64 micron. Net was washed with the water by inverting it to collect the phytoplankton attached to the net and the volume of sample was made to 100 ml. The samples were taken in separate vials and fixed in the field with 1 ml of 4 % formalin and 1 ml of Lugol's Iodine at the collection site. Ten ml of sample from each station was further concentrated by centrifuging at 2000 RPM for 10 min. For quantitative estimation of plankton, one ml of well mixed sample was taken on 'Sedgewick Rafter Cell'. To calculate density of diatoms the averages of 5 to 10 counts were made for each sample and the results are expressed as numbers of organisms per liters of sample. Qualitative study of diatoms was carried out up to the genus/species level using the standard keys given by (Edmondson, 1963; Philipose, 1960; Sarode and Kamat, 1984; APHA, 1998). Species richness of each group of plankton is considered as number of species of each group observed per visit. The number of species present in a region may be considered as its 'species richness' a frequently used measure. Species richness can be correlated positively with some measures of ecological diversity (Hurlbert, 1971). Two years (from December-2006 to November-2008) data was pooled and separated for three months and analyzed for seasonal variations, with respect to winter (December, January, February), Summer (March, April, May), Monsoon (June, July, August) and Postmonsoon (September, October, November). Mean, Standard Error of Mean (SEM) and One-Way ANOVA with No post test for various parameters for four seasons was performed. The correlation between the abiotic factors and the plankton density was calculated. The Pearson correlation was calculated by keeping diatoms as dependent variable and abiotic factors as independent variables. The P value for ANOVA is non-significant if  $P > 0.05$  (ns), Significant if  $P < 0.05$  (\*), significantly significant (\*\*\*) if  $P < 0.001$  and highly significant if  $P < 0.0001$ .

## RESULTS AND DISCUSSION

Hydrological changes are noted here with seasonal changes from dry to wet. The southwest monsoon starts by mid June and stabilizes in July and August. The stream drainage input fills the Lake, and maximal water level is in post monsoon. Water level decreases in summer due to evaporation, percolation and domestic utilization by villagers. Geologically, the area covers Deccan trap basalt rock formation and lies under monsoon semiarid deciduous forest cover.

Bacillariophyceae (Diatoms) was the most dominant family in the total phytoplankton abundance (Table.1). Density of diatoms was recorded higher in summer with  $1783 \pm 96$  ind./L,  $2035 \pm 57$  ind./L and  $2313 \pm 122$  ind./L at YLA, YLB and YLC respectively (Table.1, Fig.1) while it was decreased in monsoon and showed variations at the three stations with  $871 \pm 122$  ind./L,  $1069 \pm 172$  ind./L and  $1167 \pm 194$  ind./L respectively. The density was recorded minimal in post monsoon with non-significant differences ( $516 \pm 28$  ind./L at YLA,  $566.3 \pm 19.37$  ind./L at YLB and  $587.8 \pm 37.95$  ind./L at YLC) on all three sites investigated. The diatom densities increased in winter with insignificant differences among the three stations ( $1042 \pm 100$  ind./L at YLA,  $1054 \pm 91.96$  ind./L at YLB and  $1198 \pm 120.6$  ind./L at YLC).

**Table 1. Seasonal Variations in density (ind. /L) and species richness (no. of species) of Diatoms at YLA, YLB and YLC of Yashwant Lake during December 2006 to November 2008**

Parameters	Stations with F value	Winter	Summer	Monsoon	Postmonsoon
Diatoms Density.	YLA	1042 ± 100.8	1783 ± 96.63	871 ± 122.9	516.3 ± 28.73
	F <sub>3,20</sub> 31.77				
	YLB	1054 ± 91.96	2035 ± 57.76	1069 ± 172.6	566.3 ± 19.37
	F <sub>3,20</sub> 36.06				
Diatoms Spp. Richness	YLC	1198 ± 120.6	2313 ± 122.3	1167 ± 194.3	587.8 ± 37.95
	F <sub>3,20</sub> 30.28				
	YLA	6.33 ± 0.76	15.67 ± 1.2	12.0 ± 0.68	7.33 ± 0.95
	F <sub>3,20</sub> 22.05				
Diatoms Spp. Richness	YLB	7.33 ± 0.66	16.83 ± 0.94	14.0 ± 0.57	9.0 ± 0.57
	F <sub>3,20</sub> 38.54				
	YLC	8.0 ± 0.89	21.0 ± 0.36	15.0 ± 0.96	10.83 ± 0.87
	F <sub>3,20</sub> 48.55				




Figure 1. Seasonal variation in density of Diatoms (no. /L) at Yashwant Lake during December 2006 to November 2008

Twenty four species of diatoms (Annexure) belonging to 16 genera were recorded at Yashwant Lake. Maximal number of species were recorded in summer with  $15.6 \pm 1.2$ ,  $16.8 \pm 0.9$  and  $21.0 \pm 0.3$  at YLA, YLB and YLC respectively. Which started decreasing in monsoon with  $12 \pm 0.6$ ,  $14 \pm 0.5$  and  $15 \pm 0.9$  to postmonsoon with  $7.3 \pm 0.9$ ,  $9 \pm 0.5$  and  $10.8 \pm 0.9$  and were minimal during winter with  $6.3 \pm$

Seasonal Variation In Diatoms Density And Their Correlation .....

0.7,  $7.3 \pm 0.6$  and  $8 \pm 0.9$  at YLA, YLB and YLC respectively (Table.1, Fig.2). Both, density and species richness of Bacillariophyceae showed significant seasonal variations with  $P < 0.0001$  at all the stations.




Figure 2. Seasonal variation in species richness of Diatoms (no. of species) at Yashwant Lake during December 2006 to November 2008

Table 2. Annual variation of Physico-chemical parameters of Yashwant Lake from December 2006 to November 2008.

Parameter	YLA		YLB		YLC	
	Max.	Min.	Max.	Min.	Max.	Min.
AT °C	26.25	16.75	26.25	16.75	26.25	17.25
WT °C	23	18	26	16.2	22.7	18.7
WC %	95	60	95	60	95	60
TDS mg/l	132	99	122.5	92	138	102.5
TS mg/l	162.5	128	156	123	182	133
TSS mg/l	42.5	18.5	41	18	46	21
Trans. M	1.42	1.12	1.51	1.13	1.47	1.13
Acidity mg/l	20.5	9.25	19.5	8.25	18.75	7.5
Alkalinity mg/l	122.5	83.75	120	81.25	117.5	77.5
TH mg/l	71	48	69	49	67	46
pH	8.6	7.4	8.5	7.3	8.4	7.2
CL mg/l	33	14	30.5	13	29	12
CO <sub>2</sub> mg/l	4.8	1.3	4.8	0.2	4.9	0.4
DO mg/l	12.8	7.9	13.6	8.6	14.1	9
NO <sub>2</sub> mg/l	0.37	0.07	0.4	0.08	0.41	0.09
NO <sub>3</sub> mg/l	0.18	0.05	0.17	0.05	0.2	0.03
PO <sub>4</sub> mg/l	0.2	0.07	0.2	0.07	0.2	0.08

Table 3. Pearson correlations of Diatoms density with physicochemical parameters at YLA, YLB and YLC of Yashwant Lake during December 2006 to November 2008

Sr. No	Parameters	YLA-Site Density	YLB-Site Density	YLC-Site Density
1	Atm. Temperature	.900**	.768**	.914**
2	Water Temp.	.837**	.912**	.934**
3	Transparency	-.610**	-.536**	-.569**
4	Total Dissolved Solids	.919**	.933**	.885**
5	Total Solids	.919**	.836**	.824**
6	Total Suspended Solids	.367	.194	.230
7	Acidity	.905**	.956**	.904**
8	Alkalinity	.780**	.912**	.862**
9	pH	.890**	.919**	.909**
10	Total Hardness	.061	.217	.175
11	Chloride	.873**	.914**	.860**
12	CO <sub>2</sub>	.897**	.864**	.876**
13	DO	-.653**	-.603**	-.625**
14	NO <sub>2</sub>	.773**	.512*	.537**
15	NO <sub>3</sub>	.233	-.076	-.056
16	PO <sub>4</sub>	.856**	.736**	.783**

\*\* Correlation is significant at the 0.01 level (2-tailed).

\* Correlation is significant at the 0.05 level (2-tailed).

Phytoplankton forms the basic link of food chain between abiotic and biotic factors in the aquatic ecosystem as the metabolic activities of these organisms depend on the physicochemical factors of the aquatic environment. The quality and quantity of phytoplankton and their seasonal succession patterns have been successfully utilized to assess the quality of water and its capacity to sustain heterotrophic communities, virtually all the dynamic features of lakes such as colour, clarity, trophic state, zooplankton and fish production depend to a large degree on the phytoplankton (Goldman and Horne, 1983).

Biodiversity conservation seeks to maintain the human life support system provided by nature and the living resources essential for development. As far as water reservoirs in India are concerned three distinct plankton pulses are reported which coincide with southwest post-monsoon (September-November), winter (December-February) and summer (March-May) (Sugunan, 2000). First, the South-West Monsoon (June-August) flushing disturbs and often dislodges the standing crop of plankton. However, the destabilizing effects does not wear away immediately (as dam outlets are closed), and the allochthonous nutrient input favours some plankton growth in postmonsoon (e.g. Euglenophyceae). Second, as the post-monsoon merges into winter, the turbulence decreases and water becomes clean, the phytoplankton community progresses through a series of succession to culminate in a peak Cyanophyceae and Chlorophyceae. Third, at the end the summer plankton maxima coincide with the drastic drawdown, bringing the deep, nutrient rich areas into the fold of tropholytic zone. High temperature, bright sunlight and rapid tropholytic activities also accelerate the multiplication of phytoplankton during summer (e.g. Diatoms and Dinophyceae (Sugunan, 2000).

Diatoms are also being used increasingly as indicators of environmental changes, including studies of past climatic changes (Smol and Cumming, 2000; Wim et al, 1924 -1931). The environmental factor such as physico-chemical and biological factors influence the density and species richness of diatoms, which is reflected in their seasonal variations. Maximal diatom densities were recorded in summer at all the three stations as is also reported by earlier researches Sunkad (2002); Hujare (2005); Hulyal and Kaliwal (2009); Hafsa and Gupta (2009) and Jawale and Patil (2009). The temperature has been reported as the most important factor affecting diatom growth positively (Pearsall, 1932; Yoshitake and Imahori, 1980). In addition to temperature, George, (1961) observed that the high pH value promotes the growth of algae in general. Kamat (1965) also reported that high pH is favourable for abundant growth of diatoms. Our results are in agreement with the previous studies and it is found that For alkaline pH is favoures to flourish the members of Bacillariophyceae (Table 3). Perusal of literature indicates that Diatoms absorb large quantity of phosphates than their requirements (Rutner, 1963 and Munawar, 1970). Philipose, (1960) has reported direct relation of phosphate with diatoms. In present study, minimal to moderate phosphates were recorded in winter and summer respectively when the diatom populations were moderate to maximal. In additions, Nitrates have been given the prime

importance in diatom ecology (Patrick, 1948, Rao, 1955) and considered as the main controlling parameter in the periodicity of diatoms. However, all the three stations showed negative correlation with the diatom density with the nitrate in the present study on (Table.3). Nitrate may be utilized by diatoms. This needs further investigation in relation to other abiotic and biotic parameters.

Earlier investigation reported two peaks (i.e. in summer and winter) diatoms density George (1966); Patil (2005); Nandan and Magar (2007) and Sharma (2009) in their annual studies. In present seasonal study no such peak was observed, rather there was a steady increase in their density from winter to summer at the higher Altitudinal Lake. The effect of rains on the distribution and decline of the plankton density in general stands true for diatoms too. On the contrary, Pennak (1949, 1955) did not find any regular diatom pulse throughout the year.

Dissolved silica has a specific role in diatom growth; therefore, adequate silica supply is essential for bacillariophyceae in general. Dissolved silica is supplied to the lake by drainage water (Kobbia et al., 1992; Gad, 1992) and is also generated by remineralization within the lake. In the present study the silica is not estimated, but high density of diatoms is indicative of sufficient silica content of Yashwant Lake.

The diatoms of Yashwant Lake were represented by 24 species belonging to 16 genera which indicates availability of their distinct nutritional requirements at Yashwant Lake that favours one group over other as indicated by (Wetzel, 1990). Patrick (1973) concluded that many species of diatoms can tolerate a temperature range from 0.0 to 35°C. The study of Yashwant Lake indicates that these groups of species are found abundantly when water temperature fluctuated between  $18.75 \pm 0.3$  C of winter to  $22 \pm 0.4$  C of summer.

Palmer (1969) has listed the taxa in decreasing order of emphasis with reference to pollution index. In the present study some of these pollution tolerant species were also observed in the order of density as follows: Nitzschia, Navicula, Ulnaria, Melosira, Gomphonema, Fragilaria, Surirella, Cymbella, Pinnularia. Similar taxa were also recorded by Nandan and Mahajan, (2007) at Suki Dam, Maharashtra. Available literature indicates that Cymbella, Fragilaria species, Gomphonema are commonly found in organically rich waters while Richardson, (1968) recorded Nitzschia species is characteristic of organically rich waters. However, the clean water species of diatom such as Amphora ovalis, Cymbella sp., Pinnularia sp. were also observed in the waters of YSL. Their presence in YLA indicates that though it is not yet polluted, if care is not taken may get polluted in near future as it is having potential for deterioration and eutrophication under the influence of pollution and anthropogenic pressures.

In conclusion, present studies shows that lentic diatom communities, their temporal variation cannot be solely driven by local environmental conditions but is also determined by habitat availability in other words potential habitat is available at Yashwant Lake.

#### ANNEXURE

Diatoms of Yashwant Lake observed during December 2006 to November 2008

Bacillariophyceae (Diatoms)

1. Melosira islandica (O. Muell)
2. Ulnaria ulna (Nitz) Her. V. biceps Kuetz.
3. Ulnaria acus (Kuetz)
4. Asterionella spp
5. Frustulina spp
6. Gyrosigma accuminatum Kuetz
7. Navicula cuspidata Kuetz. V. Conspicua Venkat
8. Navicula cuspidate Kuetz. V. major Meister
9. Navicula rhynchocephala Kuetz
10. Amphora ovalis. Kuetz
11. Pinnularia interrupta W. Smith
12. Pinnularia vidarbhensis Sarode Kamat
13. Rhopalodia gibba Her O. Muell
14. Nedium longiceps Grey A. Cl. V.
15. Stauroneis obtuse Lagerst. V.
16. Surirella capronii Breb.
17. Surirella robusta Ehr.
18. Surirella sabsalsa W. Smith
19. Cymbella ventricosa Kuetz
20. Gomphonema gracile Ehr.
21. Gomphonema intricatum Kuetz
22. Fragilaria construens Ehr. Grun
23. Fragilaria zafarii Sarode Kamat
24. Nitzschia obtusa W. Smith

#### REFERENCES

- I. Wetzel, R. G. 2001. Limnology: Lakes and reservoir ecosystem (p. 1006, 3rd edn.) Academic Press Burlington.
- II. Naik, U. G. and Neelkanthan, B. 1990. Phytoplankton distribution in the Kali estuary-A seasonal study. Pollution Research, 9: 23-27.
- III. Padisak, J., Crossetti, L. O. and Naselli-Flores, L. 2009. Use and misuse in the application of the phytoplankton functional classification: a critical review with updates. Hydrobiologia, 621: 1-19.
- IV. Mooij, W. M., Hulsmann, S., De Senerpont Domis, L. N., Nolet, B. A., Bodelier, P. L. E., Boers, P. C. M., Dionisio Pires, L. M., Gons, H. J., Ibelings, B. W., Noordhuis, R., Portielje, R., Wolfstein, K. and Lammens, E. H. R. R. 2005. The impact of climate change on lakes in the Netherlands: a review. Aquatic Ecology, 39: 381-400.
- V. Pettersson, K., Grust, K., Weyhenmeyer, G. and Blenckner, T. 2003. Seasonality of chlorophyll and nutrients in Lake Erken- effects of weather conditions. Hydrobiologia, 506-509: 75-81.
- VI. Wetzel, R. G. 1990. Land-water interfaces: metabolic and limnological regulators. Verhandlungen der international Vereinigung fur theoretische und angewandte Limnologie, 24: 6-24.
- VII. Palmer, C. H. 1980. Algae and water pollution. Castle House Publication Ltd.
- VIII. Taylor, D., Dalton, C., Leira, M., Jordan, P., Chen, G., Leon-Vintro, L., Irvine, K., Bennion, H. and Nolan, T. 2006. Recent histories of six productive lakes in the Irish Ecoregion based on multiproxy paleolimnological evidence. Hydrobiologia, 571: 237-259.

- IX. Edmondson, W. T. 1963. Freshwater biology, 2nd edn. John Wiley and sons. New York, USA.
- X. Philipose, M. T. 1960. Freshwater phytoplankton of inland fisheries, Proc. Sym. Algology, ICAR., 272-291.
- XI. Sarode, P. T. and Kamat, N. D. 1984. Freshwater diatoms of Maharashtra. Saikrupa Prakashan, Aurangabad.
- XII. APHA, 1998. Standard methods for the examination of water and wastewater 20th eds. American Public Health Association. American water works Association Water Environment Federation. Washington, D.C.
- XIII. Hurlbert, S. H. 1971. The non-concept of species diversity: a critique and alternative parameters. Ecology, 52: 577-586.
- XIV. Goldman, C. R. and Horne, A. J. 1983. Limnology, Mc.Graw Hill Book Co. London, PP 464.
- XV. Sugunan, V. V. 2000. Ecology and fishery management of reservoirs in India. Hydrobiologia, 430: 121-147.
- XVI. Smol, J. P. and Cumming, B. F. 2000. Tracking long-term changes in climate using algal indicators in lake sediments. Phycology, 36: 986-1011.
- XVII. Wim, V., Elie, V., Koen, S., Koenraad, V., Mieke, S., Dominic, A. H., David, G. M., Steve, J., Bart, V. V., Vivienne, J., Roger, F., Donna, R., Victor, A. C., Cathy, K., Pieter, V. and Aaike, D. W. 2007. Historical processes constrain patterns in global Diatom diversity. Ecology, 88(8): 1924-1931.
- XVIII. Sunkad, B. N. 2002. Studies on biodiversity of wetlands with special reference to the physico-chemical factors. Ph.D. thesis, Karnatak University, Dharwad.
- XIX. Hujare, M. S. 2005. Hydrobiological studies of some water reservoirs of Hatkanangale Tahsil (Maharashtra). PhD thesis submitted to Shivaji University, Kolhapur, India.
- XX. Hulyal, S. B. and Kaliwal, B. B. 2009. Dynamics of phytoplankton in relation to physico-chemical factors of Almatti reservoir of Bijapur District, Karnataka State. Environ. Monit. Assess., 153: 45-59.
- XXI. Hafsa, S. L. and Gupta, S. 2009. Phytoplankton diversity and dynamics of Chatla floodplain lake, Barak Valley, Assam, North East India- A seasonal study. J. Environ. Biol., 30(6): 1007-1012.
- XXII. Jawale, A. K. and Patil, S. A. 2009. Physico-chemical characteristics and phytoplankton abundance of Manglore Dam, Dist. Jalgaon, Maharashtra. J. Aua. Biol., 24(1): 7-12.
- XXIII. Pearsall, W. H. 1932. Phytoplankton in the English lakes II. The composition of the phytoplankton in relation to dissolved substances. J. Ecol., 20: 241-262.
- XXIV. Yoshitake, E. and Imahori, K. 1980. Ecological studies on microalgae in lake Yunoko. Science report, 29(1): 150-158.
- XXV. George, M. G. 1961. Diurnal variations in two shallow ponds in Delhi, India. Hydrobiologia, 18(3): 265-273.
- XXVI. Kamat, M. D. 1965. Ecological notes on Kolhapur. J. Biol. Sci., 8: 47-54.
- XXVII. Rutner, F. 1963. Fundamentals of limnology, University of Toronto Press. 295 pp.
- XXVIII. Munawar, M. 1970. Limnological studies on freshwater ponds of Hyderabad, India-II. The biocenose, distribution of unicellular and colonial phytoplankton of polluted and unpolluted environment. Hydrobiologia, 41: 105-128.
- XXIX. Patrick, R. 1948. Factors affecting the distribution of diatoms Bot. Rev., 14(8): 473-524.
- XXX. Rao, C. B. 1955. On the distribution of algae in a group of six small ponds. Algal Periodicity J. Ecol., 43: 291-308.
- XXXI. George, M. G. 1966. Comparative plankton ecology of five fish ponds in Delhi, India. Hydrobiologia, 27: 81-108.
- XXXII. Patil, S. G. 2005. Hydrobiological investigations. Zool. Surv. India, Fauna of Nathasagar wetland, Wetland Ecosystem Series, 7: 7-62.
- XXXIII. Nandan, S. N. and Magar, U. R. 2007. Algal flora and physico-chemical studies of Girna Dam in Nashik District, Maharashtra. Seaweed Res. Utiln., 29(1-2): 279-285.
- XXXIV. Sharma, B. K. 2009. Composition, abundance and ecology of phytoplankton communities of Loktak Lake, Manipur, India. J. Threatened Taxa, 1(8): 401-410.
- XXXV. Pennak, R. W. 1949. Annual limnological cycles in some Colorado reservoir lakes. Ecol. Monogr., 19: 233-267.
- XXXVI. Pennak, R. W. 1955. Comparative limnology of eight Colorado mountain lakes. Uni. Colo. Stud. Biol., 2: 1-75.
- XXXVII. Kobbia, I. A., Zaki, M. Z. and Gad, Y. M. 1992. Ecology of phytoplankton in estuarine regions of Abuou Tarfaia drain in Lake Qarun (Egypt). Mansoura Sci., Bull. Egypt, 19: 141-165.
- XXXVIII. Gad, M. Y. A. 1992. Ecological studies on the algal flora of some aquatic ecosystems in EI-Fayoum district. M. Sc. Thesis, Fac. Of Sci., Cairo Univ., Giza, Egypt. Pp 222.
- XXXIX. Patrick, R. 1973. The effect of increasing light and temperature on the structure of diatom communities. Limnology Oceanography, 16: 405-421.
- XL. Palmer, C. M. 1969. A composite rating of algae tolerating organic pollution. J. Phycol., 5: 78-82.
- XLI. Nandan, S. N. and Mahajan, K. D. 2007. Studies on the diatom flora of Suki River and Suki Dam at Maharashtra. Seaweed Res. Utiln., 29(1-2): 287-298.
- XLII. Richardson, J. L. 1968. Diatoms and Lake Typology in East and central Africa. Int. Revue. Ges. Hydrobiol., 53: 299-338.



**EKHANDE A. P**

SVS's Dadasaheb Rawal College, Dondaicha.  
Dist. Dhule (M.S.) India.



**PATIL J. V**

SVS's Dadasaheb Rawal College, Dondaicha.  
Dist. Dhule (M.S.) India.

# Publish Research Article International Level Multidisciplinary Research Journal For All Subjects

Dear Sir/Mam,

We invite unpublished research paper.Summary of Research Project,Theses,Books and Books Review of publication,you will be pleased to know that our journals are

## Associated and Indexed,India

- ✍ International Scientific Journal Consortium    Scientific
- ✍ OPEN J-GATE

## Associated and Indexed,USA

- ✍ Google Scholar
- ✍ EBSCO
- ✍ DOAJ
- ✍ Index Copernicus
- ✍ Publication Index
- ✍ Academic Journal Database
- ✍ Contemporary Research Index
- ✍ Academic Paper Databse
- ✍ Digital Journals Database
- ✍ Current Index to Scholarly Journals
- ✍ Elite Scientific Journal Archive
- ✍ Directory Of Academic Resources
- ✍ Scholar Journal Index
- ✍ Recent Science Index
- ✍ Scientific Resources Database

Indian Streams Research Journal  
258/34 Raviwar Peth Solapur-413005,Maharashtra  
Contact-9595359435  
E-Mail-ayisrj@yahoo.in/ayisrj2011@gmail.com  
Website : www.isrj.net