

Vol 4 Issue 12 Jan 2015

ISSN No : 2230-7850

---

**International Multidisciplinary  
Research Journal**

*Indian Streams  
Research Journal*

---

**Executive Editor**  
Ashok Yakkaldevi

**Editor-in-Chief**  
H.N.Jagtap

---

## Welcome to ISRJ

RNI MAHMUL/2011/38595

ISSN No.2230-7850

Indian Streams Research Journal is a multidisciplinary research journal, published monthly in English, Hindi & Marathi Language. All research papers submitted to the journal will be double - blind peer reviewed referred by members of the editorial board. Readers will include investigator in universities, research institutes government and industry with research interest in the general subjects.

### *International Advisory Board*

Flávio de São Pedro Filho  
Federal University of Rondonia, Brazil

Mohammad Hailat  
Dept. of Mathematical Sciences,  
University of South Carolina Aiken

Hasan Baktir  
English Language and Literature  
Department, Kayseri

Kamani Perera  
Regional Center For Strategic Studies, Sri Lanka

Abdullah Sabbagh  
Engineering Studies, Sydney

Ghayoor Abbas Chotana  
Dept of Chemistry, Lahore University of Management Sciences[PK]

Janaki Sinnasamy  
Librarian, University of Malaya

Ecaterina Patrascu  
Spiru Haret University, Bucharest

Anna Maria Constantinovici  
AL. I. Cuza University, Romania

Romona Mihaila  
Spiru Haret University, Romania

Loredana Bosca  
Spiru Haret University, Romania

Ilie Pintea,  
Spiru Haret University, Romania

Delia Serbescu  
Spiru Haret University, Bucharest,  
Romania

Fabricio Moraes de Almeida  
Federal University of Rondonia, Brazil

Xiaohua Yang  
PhD, USA

Anurag Misra  
DBS College, Kanpur

George - Calin SERITAN  
Faculty of Philosophy and Socio-Political Sciences AL. I. Cuza University, Iasi

.....More

Titus PopPhD, Partium Christian University, Oradea,Romania

### *Editorial Board*

Pratap Vyamktrao Naikwade  
ASP College Devruk, Ratnagiri, MS India

Iresh Swami  
Ex - VC. Solapur University, Solapur

Rajendra Shendge  
Director, B.C.U.D. Solapur University,  
Solapur

R. R. Patil  
Head Geology Department Solapur  
University,Solapur

N.S. Dhaygude  
Ex. Prin. Dayanand College, Solapur

R. R. Yalikar  
Director Management Institute, Solapur

Rama Bhosale  
Prin. and Jt. Director Higher Education,  
Panvel

Narendra Kadu  
Jt. Director Higher Education, Pune

Umesh Rajderkar  
Head Humanities & Social Science  
YCMOU,Nashik

Salve R. N.  
Department of Sociology, Shivaji  
University,Kolhapur

K. M. Bhandarkar  
Praful Patel College of Education, Gondia

S. R. Pandya  
Head Education Dept. Mumbai University,  
Mumbai

Govind P. Shinde  
Bharati Vidyapeeth School of Distance  
Education Center, Navi Mumbai

G. P. Patankar  
S. D. M. Degree College, Honavar, Karnataka

Alka Darshan Shrivastava  
Shaskiya Snatkottar Mahavidyalaya, Dhar

Chakane Sanjay Dnyaneshwar  
Arts, Science & Commerce College,  
Indapur, Pune

Maj. S. Bakhtiar Choudhary  
Director, Hyderabad AP India.

Rahul Shriram Sudke  
Devi Ahilya Vishwavidyalaya, Indore

Awadhesh Kumar Shirotriya  
Secretary, Play India Play, Meerut(U.P.)

S. Parvathi Devi  
Ph.D.-University of Allahabad

S.KANNAN  
Annamalai University,TN

Address:-Ashok Yakkaldevi 258/34, Raviwar Peth, Solapur - 413 005 Maharashtra, India  
Cell : 9595 359 435, Ph No: 02172372010 Email: ayisrj@yahoo.in Website: www\_isrj.org



## DEVELOPMENT OF INDUCIBLE DRUG RESISTANCE TO AMOXICILLIN AND CEPHALEXIN IN A PROBIOTIC STRAIN OF LACTOBACILLI

Anjali A Tiwari

P.G. Department of Microbiology , Sant Gadge Baba Amravati University, Amravati .

**Abstract:**-Probiotic must be able to benefit on host through activity in human body. Lactobacillus organisms are normal inhabitants of the human intestine and vagina. They are the main ones that produce lactic acid in the digestive tract, which is important for overall health. For the prevention of intestinal disorders, probiotics can be rendered multidrug resistance to survive in the presence of antibiotics. Isolation and identification, detection and development of antibiotic resistance in this strains of lactobacillus species from commercially available probiotics and for this antibiotic susceptibility and Development of inducible drug resistance to amoxicillin and cephalexin were done and found that isolated probiotic strain was *Lactobacillus rhamnosus*. It is a Gram positive, nonmotile, non sporing, rod shaped bacteria. The size of colonies of *Lactobacillus rhamnosus* on MRS medium was of 1mm in diameter, circular, pale yellow, concave elevation, opaque and with a smooth surface and various sugar test also performed and from this result it is concluded A new strain of *Lactobacillus rhamnosus* that developed was resistant to amoxicillin and cephalexin at a concentration of 70 $\mu$ g/mL.

**Keywords:** Amoxicillin And Cephalexin , Probiotic Strain ,Lactobacillus .

### INTRODUCTION

Probiotics are the live microorganism which when administered in adequate amount confirms health benefit, to host and maintain intestinal flora. Infectious diseases are the biggest problem in human being and every year gastrointestinal infections are responsible for significant morbidity and mortality worldwide (Culligan *et al*, 2009). World Health Organization (WHO, 2004) estimates there to be more than four billion episodes of diarrhoeal disease annually, while there were 2.2 million deaths attributable to enteric infection, making it the fifth leading cause of death at all ages worldwide. Probiotics can be used in treatment and prevention of enteric infection and chronic inflammatory disorders of the gastrointestinal tract (Gill and Guarner, 2004). The following are microorganisms considered to be human Probiotics:

**Lactobacillus species:** *L. acidophilus*, *L. amylovorus*, *L. brevis*, *L. casei*, *L. casei* subsp. *Bifidobacterium species*: *B. adolescentis*, *B. bifidum*, *B. breve*, *B. infantis*, *B. lactis* (*B. animalis*).

**Other lactic acid bacteria:** *Enterococcus faecium*, *Lactococcus lactis*, *Leuconostoc mesenteroides*,

**Nonlactic acid bacteria:** *Bacillus subtilis*, *Escherichia coli* strain nissle, *Saccharomyces boulardii*,

Lactobacilli constitute a major part of the natural microflora of human intestine. These probiotic organisms when present in sufficient number can create a healthy equilibrium between beneficial and potentially harmful microflora in gut by creating unfavourable condition for the growth of commonly occurring intestinal pathogen (Tagg and McGiven, 1971; Salminen *et al.*, 1998).

Few *Lactobacillus* strains are able to inhibit the growth and adhesion of *Candida albicans* or other Candida species, and there is no solid evidence to indicate that intravaginal administration of lactobacilli can eradicate yeast infection. However, there is some evidence to suggest that lactobacilli ingestion and vaginal use can reduce the risk

Anjali A Tiwari , "DEVELOPMENT OF INDUCIBLE DRUG RESISTANCE TO AMOXICILLIN AND CEPHALEXIN IN A PROBIOTIC STRAIN OF LACTOBACILLI" Indian Streams Research Journal | Volume 4 | Issue 12 | Jan 2015 | Online & Print

## **Development Of Inducible Drug Resistance To Amoxicillin And Cephalexin In A Probiotic Strain Of Lactobacilli**

---

of recurrences (Hilton et al, 1992; 1995).

### **AIMS AND OBJECTIVES**

For the prevention of intestinal disorders, probiotics can be rendered multidrug resistance to survive in the presence of antibiotics. Therefore the present study aims at,

- ❖ Isolation and identification of the strains of lactobacillus species from commercially available probiotics.
- ❖ Detection of the antibiotic sensitivity in the isolated lactobacillus strain.
- ❖ Development of antibiotic resistance in this strain by inducible drug resistance.

### **MATERIALS AND METHODS**

Materials: The commercially available probiotic products like Ecobion, Vibact, Bioclin, etc were procured from the local medical stores.

Media (Hi media make)

1. Peptone physiological solution.
2. Phosphate buffer
3. De Man Rogosa and Sharp agar Composition
4. Antibiotics (Sigma) Amoxicillin, Cephalexin

#### **A. Commercial probiotics and their contents.**

**1.Biocline -***Lactobacillus acidophilus, Lactobacillus rhamnosus, Bifidobacterium longum Lactobacillus casei*

**2.Ecobion-** *Lactobacillus acidophilus, Lactobacillus rhamnosus, Bifidobacterium bifidum, Bifidobacterium longum, Saccharomyces boulardii, Streptococcus thermophilus*

**3.Vibact -***Streptococcus faecalis T-110JPC, Clostridium butyricum TO-A, Bacillus mesentericus TO-AJPC, Lactic acid bacteria (Lactobacillus sporogenes )*

**5.Antibiotic discs** (Hi media, Mumbai Ltd.)

Names of antibiotics	Concentrations
Amikacin	30 µg
Amoxycillin	10 µg
Ampicillin	10 µg
Ceftazidime	30 µg
Ceftriaxome	30 µg
Cephalexin	30 µg
Cephaxitin	30 µg
Chloramphenicol	30 µg
Cloxacillin	5 µg
Co-trimoxazole	25 µg
Erythromycin	15 µg
Gentamicin	10 µg
Penicillin	25 µg
Penicillin-v	3 µg
Piperacillin	100 µg
Tetracycline	10 µg

## METHODS

### 1)Isolation of probiotic strain from commercial product

Commercially available powdered probiotic was inoculated in peptone physiological solution. 100mg sample was suspended in 100 mL of peptone physiological solution and incubated at 37°C for 24h. After incubation loopful culture from broth was inoculated on MRS agar medium and again incubated at 37°C for 24–48h. The colonies were identified by cultural and morphological characteristics and sugar fermentation tests.. Sugar fermentation test was done by using disc of sugars. Phenol red base broth was used in which the sugar disc were added and then the culture was inoculated into the broth. This was incubated at 37°C for 24h. Change in colour from red to yellow indicated sugar fermentation with formation of acid.

### 2)Antibiotic susceptibility test

Antibiotic susceptibility test was performed by Kirby Bauer disc diffusion method. The antibiotics to which the probiotic strain was highly sensitive were selected for further study.

### 3)Development of inducible drug resistance to amoxicillin and cephalexin.

#### a. Preparation of antibiotic solution

Pure antibiotic in powder form was procured from Sigma Company. Phosphate buffer of pH 6.0 was used as a solvent as well as diluent for amoxicillin and cephalexin. Stock solution of antibiotics was prepared by adding 500mg of antibiotic powder in 100mL of sterile phosphate buffer this was the stock solution and the concentration was 5mg/mL. Working solution was prepared by diluting 10 ml of stock solution in 100 mL of sterile phosphate buffer. This was first working solution and the concentration of this solution was 500µg/mL. From this first working solution 10 mL of solution was again diluted to 100 mL of phosphate buffer. This was second working solution and the concentration was 50µg/mL. Again from this solution 10 mL solution was taken to make the solution having concentration 5mg/mL. In similar manner dilution was carried out to get the final concentration of 5µg/mL. The prepared antibiotic solution was added aseptically to sterile molten nutrient agar in increasing order and allowed to solidify.

#### b.Method for developing inducible drug resistance.

The isolated probiotic strain was inoculated on the nutrient agar plate containing 5µg/mL of antibiotic and incubated at 37°C for 24 h. after incubation growth observed was transferred to nutrient agar containing 10µg/mL of antibiotic and again incubated. In such a way the growth was transferred to nutrient agar medium containing increasing concentration of antibiotics.


In a similar manner serial passage of the strain was carried out till 70µg/mL concentration of antibiotics. Simultaneously the initially isolated probiotic strain was also inoculated on the plate with increasing concentration of antibiotics to detect the development of drug resistance. The probiotic strain that resisted to 70µg/mL of antibiotic was again transferred to plane nutrient agar / MRS agar medium and then they again subcultured on nutrient agar with 70µg/mL of antibiotic to determine the stability of the acquired drug resistance.

## RESULTS

From the three products different probiotic isolates were obtained. Among them one isolate was selected and identified by morphological, cultural characteristics and biochemical tests.The isolated probiotic strain was *Lactobacillus rhamnosus*. It is a Gram positive, nonmotile, non sporing, rod shaped bacteria. The size of colonies of *Lactobacillus rhamnosus* on MRS medium was of 1mm in diameter, circular, pale yellow, concave elevation, opaque and with a smooth surface. Sugar fermentation test was carried out to confirm the isolate. The isolated *Lactobacillus rhamnosus* was sensitive to all the antibiotics tested.

**Development Of Inducible Drug Resistance To Amoxicillin And Cephalexin In A Probiotic Strain Of Lactobacilli**

*L. rhamnosus* isolated from probiotic product inoculated on media with antibiotic



## DISCUSSION

The present investigation was aimed at isolation, identification, antibiotic susceptibility testing and

#### **Development Of Inducible Drug Resistance To Amoxicillin And Cephalexin In A Probiotic Strain Of Lactobacilli**

---

development of inducible antibiotic resistance in a strain of probiotic from a commercial product. Accordingly a strain of *Lactobacillus rhamnosus* was isolated from a commercially available probiotic powder. It was identified by cultural and morphological characteristics and confirmed by sugar fermentation test. Similar isolation and characterization of probiotic bacteria from commercial product was carried out by R Temmerman. They processed 55 european probiotic products to identify and the antibiotic resistances of the bacterial isolates recover from this product and found that from 30 dried food supplement and 25 dairy products yielded a total 268 bacterial isolates.

Resistance to amoxicillin and cephalexin was induce in the strain of *Lactobacillus rhamnosus* was developed that could resist the concentration of amoxicillin cephalexin upto 70 $\mu$ g/mL. If such probiotics strains are consumed probiotics as the probiotic will not be destroyed by the antibiotics.

According to the study conducted by Behira Belkacem and Kihal Mebrouk, on antibiotic resistance of some lactobacilli isolated from the gut microflora of broiler such type of antibiotic resistance could also be induced in the gut when both the antibiotics and probiotics are consumed together.

#### **CONCLUSION**

In the present study *Lactobacillus rhamnosus* was isolated and identified from commercially available probiotic products. Its antibiotic susceptibility was determined to various antibiotics. It was found to be completely sensitive to all the antibiotic tested. While the resistant to amoxicillin and cephalexin was induced in *Lactobacillus rhamnosus*, it readily get resistance to the two drugs. It could resist upto 70 $\mu$ g/mL of the drug. This indicates that the bacteria can very easily acquired resistance to antibiotics. A new strain of *Lactobacillus rhamnosus* that developed was resistant to amoxicillin and cephalexin at a concentration of 70 $\mu$ g/mL. Thus it can be conclude from the study that such type of inducible drug resistance can be developed in probiotics so that it could be effectively administered during the course of antibiotics and prevent the antibiotic induced diarrhoea.

#### **REFERENCES**

- 1.Arunachalam K, Gill HS and Chandra RK, (2000). Enhancement of natural immunity function by dietary consumption of *Bifidobacterium lactis* HN019. European J Clin Nutr, 54: 1-4.
- 2.Collins JK, Thornton G and O'Sullivan GO, (1998). Selection of probiotic strains for human applications. Int Dairy J 8: 487-490.
- 3.Culligan EP, Hill C, and Sleator RD, (2009). Probiotics and gastrointestinal disease: successes, problems, and future prospects. Gut pathogens, 1(19): 1-12.
- 4.Gill H S, and Guarner F, (2004). Probiotics and human health: a clinical perspective. Postgrad Med J, 80: 516-526.
- 5.Gill, H.S. and Guarner F., (2004). Probiotics and human health: a clinical perspective. Postgrad Med J, 80: 516-526.
- 6.Hilton E, Isenberg HD, Alperstein P, France K and Borenstein MT, (1992). Ingestion of yogurt containing *Lactobacillus acidophilus* as prophylaxis for candidal vaginitis. Ann Intern Med, 116: 353-357
- 7.Hilton E, Rindos P and Isenberg HD (1995): *Lactobacillus GG* vaginal suppositories and vaginitis. J Clin Microbiol, 33: 1433.
- 8.Saavedra JM, Bauman NA, Oung I, Perman JA and Yolken RH, (1994). Feeding of *Bifidobacterium bifidum* and *Streptococcus thermophilus* to infants in hospital for prevention of diarrhoea and shedding of rotavirus. Lancet, 344: 1046-9.
- 9.Tagg , J.R. and McGiven,(1971). Assay system for bacteriocines. Appl Microbial, 21,943.
- 10.World Health Organization, (2004). <http://www.who.int/info/healthinfo/globalburden/ease/GBD report part 2.pdf>.

# **Publish Research Article International Level Multidisciplinary Research Journal For All Subjects**

Dear Sir/Mam,

We invite unpublished Research Paper,Summary of Research Project,Theses,Books and Book Review for publication,you will be pleased to know that our journals are

## **Associated and Indexed,India**

- \* International Scientific Journal Consortium
- \* OPEN J-GATE

## **Associated and Indexed,USA**

- Google Scholar
- EBSCO
- DOAJ
- Index Copernicus
- Publication Index
- Academic Journal Database
- Contemporary Research Index
- Academic Paper Databse
- Digital Journals Database
- Current Index to Scholarly Journals
- Elite Scientific Journal Archive
- Directory Of Academic Resources
- Scholar Journal Index
- Recent Science Index
- Scientific Resources Database
- Directory Of Research Journal Indexing

Indian Streams Research Journal  
258/34 Raviwar Peth Solapur-413005,Maharashtra  
Contact-9595359435  
E-Mail-ayisrj@yahoo.in/ayisrj2011@gmail.com  
Website : [www.isrj.org](http://www.isrj.org)